

List of Figures

Figure 1.1: The left-hand palm is non-superimposable on the mirror image by any set of rotations and translations.	2
Figure 1.2: Schematic diagram of chiral molecule. Image formed on the mirror is non-superimposable mirror image.	2
Figure 1.3: Types of Stereoisomers with example.....	4
Figure 1.4: Unpolarized light travels through a polarizer, which converts it into PPL.	5
Figure 1.5: Horizontal and Vertical Polarization representation.	6
Figure 1.6: (a) Schematic representation of circularly polarized light. (b) Schematic of right and left circularly polarized light.	7
Figure 1.7: (a) Photograph of Louis Pasteur (Gal, 2011) (b) Enantiomeric crystals of tartaric acid separated by Louis Pasteur.(<i>Tartaric Acid</i>).	8
Figure 1.8: (a) Photograph of Jacobus H. van't Hoff (Krätz, 1974) (b) Joseph A. Le Bel (Encyclopaedia, 2023)	9
Figure 1.9: An example of a chiral molecule (bromo-chloro-fluro methane) where the central carbon atom is the chiral center attached to four different groups. It exists in two stereoisomeric forms, both mirror images of each other.	10
Figure 1.10: (a) Stereoisomeric structures of tartaric acid. D-tartaric acid and L-tartaric acid (optically active) and meso Tartaric acid (optically inactive). (b) Chiral allenes have two different groups attached on terminal carbon atoms.	11
Figure 1.11: (a) Photograph of Kurt Mislow (Buda et al., 1992) (b) Jay Siegel (S.Siegel, 2009)	12
Figure 1.12: Bromo-chloro-fluro-methane has one chiral center, while 2-bromo-3-chlorobutane has two chiral centers.	13
Figure 1.13: Enantiomers of helical chiral molecule.(Rice et al., 2018)	13
Figure 1.14: Planar chiral Ferrocenes. (Zhu et al., 2016)	14
Figure 1.15: Enantiomers of axially chiral molecule BINOL.(Krajnc & Niemeyer, 2022)	14
Figure 1.16: The rotation of plane polarized light through dextrorotatory enantiomer is clockwise, levorotatory is anticlockwise, and optical rotation is cancelled in racemic mixture.	15
Figure 1.17: Rotation of plane of polarization through optically active sample.	16
Figure 1.18: Rotation of plane of polarization due to difference in absorption of RCP and LCP.	17
Figure 1.19: Successive alternate right-handed and left-handed quartz prisms geometry that separates RCP and LCP.	18

Figure 1.20: Schematic and working of polarimeter.	21
Figure 1.21: Schematic of half shade plate used in Laurent’s polarimeter.....	22
Figure 1.22: (a) Schematic of biquartz plate used in biquartz polarimeter. (b) Colours observed through biquartz plate.....	24
Figure 1.23: Schematic of Lippich polarimeter.	24
Figure 1.24: Schematic of circular dichroism spectropolarimeter. The differential absorption of RCP and LCP through a chiral sample helps to investigate structure of a molecule.	26
Figure 1.25: Schematic diagram of the Rayleigh (elastic scattering) and Raman scattering (stokes and anti-stokes inelastic scattering).	28
Figure 1.26: Polarizability ellipsoid.(Vasilescu & Adrian-Scotto, 2010).....	29
Figure 1.27: Polarizability ellipsoid of water molecule for different vibrational modes.....	29
Figure 1.28: Types of Vibrational optical activity (VOA).	31
Figure 1.29: Energy level diagrams of different forms of ROA.(Laurence A Nafie et al., 1991)	32
Figure 1.30: Vibrational circular dichroism (VCD) measures difference in absorbance of right and left circularly polarized light.	34
Figure 1.31: Negative refractive index metamaterials bend light in the opposite direction compared to conventional positive refractive index materials.	35
Figure 2.1: Photodetector and Detector output measurement unit.	40
Figure 2.2: Prism Table Spectrometer	41
Figure 2.3: Analytical Balance	41
Figure 2.4: Schematic of Abbe Refractometer.	42
Figure 2.5:(a) Newton's Rings Apparatus, (b) Interference rings formed with a monochromatic source (sodium light)	45
Figure 2.6: Schematic showing geometrical details of plano convex lens and plane glass plate.....	46
Figure 2.7: Schematic of Circular Dichroism Spectroscopy.	48
Figure 2.8 (a) Syntek symmetric diamond anvil cell (DAC), (b) Schematic arrangement of pair of diamonds, inside the cell.	49
Figure 2.9: Pressure calibration with ruby fluorescence.....	50
Figure 2.10: Stereo Zoom Microscope	50
Figure 2.11:(a) Raman Spectrometer, (b) Raman Spectrometer Schematic representation. ...	52
Figure 2.12: (a) Schematic representation of Raman optical activity setup. (b) Modified Raman Optical Activity setup.....	54

Figure 2.13: Transmission of IR radiation through fast and slow axis.....	55
Figure 3.1: The calcite crystal's optical axis and blunt corner are displayed. The blunt corner (Corner A) of the anisotropic crystal is the corner where the surface planes intersect to make three obtuse angles with their edges, and the dotted line going through it is the crystal's optic axis.....	59
Figure 3.2: Propagation of e-ray and o-ray in a negative uniaxial crystal is depicted schematically. (a) The incident ray at any arbitrary angle with the crystal's optic axis, (b) Incident ray perpendicular to the crystal's optic axis or (c) Incident ray parallel to the crystal's optic axis.....	61
Figure 3.3: Optical indicatrix for (a) Isotropic, (b) Uniaxial positive, and (c) Uniaxial negative crystals are shown. Isotropic crystals have same refractive indices in all directions, and when it comes to uniaxial crystals, the refractive indices vary between two extreme values and form uniaxial prolate and uniaxial oblate indicatrix.....	62
Figure 3.4: A biaxial positive and biaxial negative crystal optical indicatrix is shown. Biaxial crystals have one intermediate refractive index (β) that lie between two extreme refractive indices (α and γ).	63
Figure 3.5: A flowchart illustrating how an optical medium is classified according to its light-propagation characteristics.....	64
Figure 3.6: Light splits into fast and slow rays when it enters an anisotropic crystal with a thickness of 'd'. The slow ray is represented by a blue wave in the crystal, whereas the fast ray is represented by a brown wave. In comparison to the fast ray, the slow ray covers a shorter distance and develops retardation ' Δ '.	65
Figure 3.7: Plane of polarization rotated through an angle $2\theta^\circ$ by half wave plate.....	66
Figure 3.8: A quarter wave plate's (QWP) response to linearly polarized light is depicted visually. When optic axis of QWP is parallel to the polarizer transmission axis no change in polarization but when it is oriented at 45° , it changes the state of polarization and circularly polarized is generated.	68
Figure 3.9: Generation of right/left circularly polarized light when fast axis of quarter wave plate is oriented at $+45^\circ$ and -45° . (Figures are drawn by observing optical geometry from the outside).....	68
Figure 3.10: (a) A two-beam Shimadzu 460 infrared spectrometer schematic arrangement is shown. The transmittance spectra are recorded and detected by rotating the fast and slow axes of the quartz and mica wave plates. Maximum (minimum) transmittance values are obtained by keeping the fast axis (Figure 10a/(slow axis) Figure 3.10 (b) parallel to the polarizer and analyzer axes. (Guohua et al., 1990).....	69
Figure 3.11: A quarter wave plate with a known axis is used to determine fast axis of a quarter wave plate whose fast axis is unknown. (Galgano & Henriques, 2005)	70
Figure 3.12: Schematic arrangement shows how metal's ability to modify phase is used to determine fast axis of a quarter wave plate. The transmission axes of two linear polarizers are	

crossed at 45° to the left and right of the vertical direction, respectively. (Galgano & Henriques, 2005; Logofatu, 2002)	71
Figure 3.13: An overview of the setup used by M.H. Chiu et al. to calculate the fast axis and phase retardation using heterodyne interferometry. (BS- beam splitter; M- Mirror; P- Polarizer; EOM- Electric optic modulator; A _t and A _r -analyzer; W- wave plate; D _t and D _r ,- photodetector. (Chiu et al., 1996)	72
Figure 3.14: Destructive interference is observed as resolved components from an anisotropic crystal are cancelled along the transmission direction of analyzer.....	76
Figure 3.15: Constructive interference is observed as resolved components from an anisotropic crystal are added and allowed to pass through the analyzer.	76
Figure 3.16: When monochromatic light is allowed to pass through the quartz wedge placed between the crossed polarizer and analyzer. Bright and dark bands appear due to interference of resolved components of fast and slow rays.	77
Figure 3.17: Michel Levy interference colour chart (Adapted with permission from “Molecular Expressions at Florida State University” (Robinson & Davidson)	80
Figure 3.18:(a) Schematic of the experiment's setup (b) Our experimental actual setup. The light source is a mercury vapour lamp. Axes are represented as straight lines to indicate how each optical device is oriented.	81
Figure 3.19: Figures depict colour as they appear when a quarter-wave plate composed of mica is tilted around one of the axes and seen through the telescope of the spectrometer. On the left, rotation around the fast axis causes white light to turn grey, and on the right, rotation around the slow axis causes white light to turn yellow.	83
Figure 3.20: Figure depicts the colours as they appear through the telescope eyepiece of the spectrometer when a quarter-wave plate made of quartz is tilted around one of the axes. Due to rotation on the fast axis (reduced retardation), on the left, the light-yellow colour changes to cyan green, while on the right, the pale-yellow colour changes to pink (more retardation).	83
Figure 3.21:The left plot relates to photocurrent fluctuation with rotation angle caused by Mercury source, whereas the right plot corresponds to photocurrent variation caused by Sodium source when quarter mica wave plate is placed between cross polarizers. The graph's peaks represent the angles when the intensity is at its maximum. At maximal intensity, the fast and slow axes' positions may be determined.	84
Figure 3.22: A quarter wave plate is placed between a polarizer and an analyzer in a parallel arrangement and rotated to monitor intensity fluctuations.	85
Figure 3.23: QWP made up of mica is placed between parallel polarizers and rotated, and intensity variation as a function of rotation is plotted. The left graph corresponds to a Mercury source, and the right graph corresponds to a Sodium source.	86
Figure 4.1: Plane polarized light is the vector sum of two oppositely rotating left and right circularly polarized light (a) No optical rotation when plane polarized light is incident in	

chiral sample. (b) As plane polarized light exits from chiral sample its plane of polarization is rotated.	89
Figure 4.2: Schematic arrangement of Newton’s Rings Setup.	94
Figure 4.3: Newton Rings formed with monochromatic sodium light source.	95
Figure 4.4: Modified Newton’s Ring Setup, polarized light falls on the cavity chamber.	96
Figure 4.5: Schematic of modified Newton’s rings setup. Circularly polarized light is generated by placing quarter wave plate in the path.	97
Figure 4.6 Plot corresponds to Refractive index of both enantiomers of Glucose, Mannose, Tartaric acid, BINOL, Alpha methyl benzyl amine and Alpha ethyl benzyl amine with respect to different polarizing angles in the interval of 30°.	99
Figure 5.1: Chemical structure of Na ₂ S ₂ O ₄	111
Figure 5.2: Sodium Dithionite’s (Na ₂ S ₂ O ₄) ambient Raman spectrum.	111
Figure 5.3: Chirality induction by Na ₂ S ₂ O ₄ : Hypothetical Reaction Scheme.	112
Figure 5.4: Sodium dithionite and acetophenone mixed well in a mortar and pestle instantly weaken the S-S stretching mode. A drop of methanol reduces the intensity of the S-S stretching bond.	113
Figure 5.5: Sodium dithionite and acetophenone mixed well in a mortar and pestle instantly weaken the carbonyl bond frequency, and after adding a drop of methanol, break the carbonyl bond completely.	113
Figure 5.6: Sodium dithionite and benzil are mixed well in a mortar and pestle. It instantly weakens the S-S stretching bond, and after adding a drop of methanol, it weakens the carbonyl bond.	114
Figure 5.7: Raman spectra of Sodium dithionite (Na ₂ S ₂ O ₄) under quasi hydrostatic pressure (0-34.5GPa).(Shah et al., 2022)	118
Figure 5.8: Raman modes' pressure dependency under quasi-hydrostatic pressure. The filled and open symbols represent compression and decompression, respectively.	119
Figure 5.9: Under quasi-hydrostatic pressure, the normalized Raman spectra of Na ₂ S ₂ O ₄ – Black (At lowest pressure-0 GPa) and Red (fully decompressed).	120
Figure 5.10: Raman spectra of Sodium dithionite (Na ₂ S ₂ O ₄) under non-hydrostatic pressure (0-30.2 GPa).(Shah et al., 2022)	121
Figure 5.11: Raman modes' pressure dependency under non-hydrostatic pressure. The filled and open symbols represent compression and decompression, respectively.	122
Figure 5.12: Under non-hydrostatic pressure, the normalized Raman spectra of Na ₂ S ₂ O ₄ – Black (At lowest pressure-0 .34 GPa) and Red (fully decompressed).	123
Figure 5.13: Structure of crystalline sodium dithionite (Na ₂ S ₂ O ₄).	124
Figure 6.1: Different types of Raman Optical Activity. (Nafie et al., 1991)	131

Figure 6.2: Chemical structure of R BINAP and S BINAP.....	133
Figure 6.3: Modified Raman setup to record Raman optical activity (ROA) of chiral samples. QWP stands for quarter wave plate.	134
Figure 6.4: Schematic of structural diagram of BINOL with numbering of atoms.....	135
Figure 6.5: Normalized ambient Raman spectra of R BINAP (black line) and S BINAP (red line).	136
Figure 6.6: Ambient Raman Optical Activity (ROA) spectra of R BINAP (black line) and S BINAP (red line).....	139
Figure 6.7: Raman Optical activity (ROA) spectra of R BINAP at 0 GPa and 0.08 GPa.	141
Figure 6.8: Raman Optical activity (ROA) observed stretching mode $\sim 1378\text{ cm}^{-1}$ R BINAP with compression.	141
Figure 6.9: Raman Optical activity (ROA) spectra of S BINAP at 0 GPa and 0.08 GPa.	143
Figure 6.10: Raman Optical activity (ROA) spectra of S BINAP with compression.....	144
Figure 6.11: Raman spectra of R BINAP (2, 2'- bis (diphenyl phosphino)- 1,1' binaphthyl) at non-hydrostatic pressure (0-11.77GPa).	146
Figure 6.12: Raman spectra of solid R BINAP. (fully decompressed).....	147
Figure 6.13: Pressure dependence of Raman modes under non-hydrostatic pressure. The filled symbols denote compression and open symbols denote decompression. In ω vs P plot, Sp and Re labels are used to indicate splitted peaks, and reversible peaks.	148
Figure 6.14: Raman spectra of S BINAP (2, 2'- bis (diphenyl phosphino)- 1,1' binaphthyl) at non-hydrostatic pressure (0-12.62GPa).	150
Figure 6.15: Decompression stack plot of S BINAP.	151
Figure 6.16: Pressure dependence of Raman modes under non-hydrostatic pressure for S BINAP. The filled symbols denote compression and open symbols denote decompression. Sp and Re labels are used to indicate splitted and reversible peaks.....	152
Figure 8.1: The overlapping Ruby fluorescence spectra of Quasi hydrostatic (BLACK) and non hydrostatic (RED) pressure environments observed at different pressures.	160
Figure 8.2: Ruby fluorescence decompression stack plot in quasi hydrostatic pressure environment is shown. Ruby peaks are completely reversible with decompression.	161
Figure 8.3: Ruby fluorescence decompression stack plot in non hydrostatic pressure environment is shown.	161
Figure 8.4: Powder X-Ray Diffraction pattern of Sodium dithionite $\text{Na}_2\text{S}_2\text{O}_4$	164
Figure 8.5: X- ray diffractogram of all observed phases are shown. Red (sodium dithionite), green (sodium sulfite), Dark red (β -Sulfur), Blue (sulphur), and Pink (sodium oxide).	167
Figure 9.1: Refraction of light traveling through two different media.	209

Figure 9.2: In a ray diagram, two different bending directions of light are represented. Blue refracted ray represents normal (positive) refraction through DPM, and the purple refracted ray represents refraction through NIM.	211
Figure 9.3: An array of continuous wires and split ring resonators.....	212
Figure 9.4: Schematic of an LC oscillator.	213
Figure 9.5: Schematic of SRR working as an LC oscillator.	215
Figure 9.6: In solids, the positive charge is fixed in the core and is surrounded by free, negatively charged electrons.....	216
Figure 9.7: Pendry et al. presented a model consisting of conducting metallic cylinders to achieve negative permeability. (J B Pendry et al., 1999).....	218
Figure 9.8: In Model B, conducting cylinders are made up of split ring structures separated by distance d	219
Figure 9.9: Model C, a magnetic field is applied parallel to an N -turn coil.	219
Figure 9.10: Split ring resonator parameters constructed by Smith et al.....	220
Figure 9.11: Propagating waves oscillate as Sine waves and evanescent waves decay exponentially.....	221
Figure 9.12: Image formation in normal lens.	221
Figure 9.13: (a) Negative refractive index metamaterial (NIM) lenses diverge propagating waves from a point source before converging to form images. (b) Evanescent waves are strengthened in negative refractive index metamaterials and contribute significantly to image formation.....	222
Figure 9.14: Schematic of an invisibility cloak that hides objects from view.	223
Figure 9.15: (a) A NIM beam tilting antenna installed between two dielectric resonator antennas. (b) Schematic of NIM unit cell.	223
Figure 9.16: (a) Schematic of beam tilting terahertz antenna with NIM. (b) NIM embedded with graphene.....	224
Figure 9.17: Double refraction occurs in chiral nihility material. Both refracted rays are circularly polarized (CP); one circularly polarized light is refracted positively, and the other is refracted negatively.....	225
Figure 9.18: Asymmetric bilayer cross wires fabricated on a copper clad FR-4 board show strong chirality and act as chiral metamaterials.	226
Figure 9.19: Schematic of chiral metamaterial made from an insulated strip of metal.	227

List of Tables

Table 2.1. Chemicals Details	37
Table 4.1: List of samples studied and sample concentration	93
Table 4.2: Refractive index comparison study for both samples with the particular CP light.	100
Table 4.3: Refractive index comparison for particular sample with both CP lights.....	102
Table 5.1: Raman modes at ambient condition of crystalline Sodium Dithionite ($\text{Na}_2\text{S}_2\text{O}_4$).	115
Table 6.1: Raman modes at ambient condition of solid 1,1'- Bi-2-naphthol (BINOL) excited at 514.5 nm. ν , bond stretching; τ , out-of-plane (naphthyl) torsion of carbon atoms; τ_{butt} , butterfly torsion between two naphthyl rings; γ , out-of-plane (naphthyl) wagging of hydrogen atoms, δ , in-plane (naphthyl) bond bending.	137
Table 8.1: Experimental details of X-Ray Diffraction.....	162
Table 8.2: The intensity of diffracted X-rays with respect to the incident angle.....	165
Table 9.1: Materials classified on the basis of ϵ and μ values.	210